

**PERFORMANCES
DIFFERENCE ON CLIMATE BETWEEN
AR4 AND AR5**

MUHAMMAD IZHAR BIN ISMAIL

B. ENG(HONS.) CIVIL ENGINEERING

UNIVERSITI MALAYSIA PAHANG



STUDENT'S DECLARATION

I hereby declare that the work in this thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at Universiti Malaysia Pahang or any other institutions.

(Student's Signature)

Full Name : MUHAMMAD IZHAR BIN ISMAIL

ID Number : AA15250

Date : 30 MAY 2019

PERFORMANCES DIFFERENCE ON CLIMATE BETWEEN AR4 & AR5

MUHAMMAD IZHAR BIN ISMAIL

Thesis submitted in fulfillment of the requirements
for the award of the
B.Eng (Hons.) Civil Engineering

Faculty of Civil Engineering and Earth Resources
UNIVERSITI MALAYSIA PAHANG

MAY 2019

ACKNOWLEDGEMENTS

I would like to express my sincere gratitude towards my supervisor Dr Nurul Nadrah Aqilah Binti Tukimat for her guidance, ideas, continuous encouragement and support in making this research possible. She was helpful in many ways in my research, from bright ideas towards consulting on how to approach problems to solve them. Also, she gave me an insight towards what should be called life-long learning experience. She had strict principles and a firm belief on what she was doing. I really appreciate the help that she has given to me and my teammates in completing our study. Without her progressive thinking and progressive attitude, I would never made it towards the end of the Bachelor's Degree I'm currently pursuing.

A special gratitude I would like to forward towards my family especially my mum, Normah for their support through my ups and downs, making sure that they have my back whenever I needed them and plus, encouraging me to go further and surpass myself in all kinds of aspects. They're also my backbone towards me finishing this study and helped me in understanding, having patience and finally, being the ones I can turn back to if I am in need of help.

My sincere thanks towards my teammates, Fatin & Basri in helping me finish this research and the cooperation in making the study more understandable, viable to simulate and support in me finishing the research.

I would like to acknowledge my housemates, Banglo 2 Tingkat inhabitants especially Ashrawi, Asyraf, Iswandy, Kumpulan Muzik UMP, including En Fakhruddin, En Rani, Izzat, Fazrul, Firdhaus for their tremendous support and confidence in me towards concluding my life here in UMP. I also would like to acknowledge their suggestions and comments, which was crucial for in completing the study.

.

ABSTRAK

Intergovernmental Panel on Climate Change (IPCC) merevisi kesan gas rumah kaca (GHG) ke dalam sistem iklim dan mengeluarkan Laporan Penilaian Kelima (AR5) pada tahun 2014. Dengan AR5, perubahan iklim telah diklasifikasikan berdasarkan tahap radiasi memaksa dikenali sebagai Laluan Konsentrasi Perwakilan (RCP2.6, RCP4.5, dan RCP8.5). Versi ini sedikit berbeza dengan versi AR4 yang berdasarkan kumpulan GHG yang dikenali sebagai A1, B1, A2, dan B2. Pengubahsuaian dalam penilaian perubahan iklim akan menjejaskan ketepatan unjuran iklim dalam jangka panjang. Oleh itu, matlamat utama kajian ini adalah untuk menentukan kesan dalam penilaian perubahan iklim antara Laporan Penilaian Keempat (AR4) dengan AR5. Kajian itu difokuskan pada Malaysia Timur termasuk Kelantan dan Terengganu. Dalam kajian ini, Model Downscaling Statistik (SDSM) digunakan sebagai model iklim statistik untuk menilai perbezaan persembahan iklim. Sementara itu, model peredaran umum (GCM) yang disediakan oleh Pemodelan dan Analisis Iklim (CanESM2) digunakan untuk penjanaan iklim jangka panjang. Merujuk kepada hasil, ramalan p-u, r500, dan r850 adalah pembolehubah yang paling mempengaruhi dalam membentuk suhu tempatan dan hujan di kawasan. Ketepatan penjanaan iklim dikawal oleh % MAE yang lebih rendah dengan Korelasi tinggi (R) dalam keputusan yang dikalibrasi dan disahkan. Simulasi suhu berjaya menghasilkan 0.6% MAE dengan R hampir 1.0. Sementara itu hujan di Terengganu dan Kelantan dihasilkan kurang daripada 14% MAE dengan 0.99 R. Berdasarkan perbandingan prestasi antara GCM dan data sejarah, RCP4.5 (AR5) dan SRES A2 (AR4) telah dipilih sebagai tahap pemantauan radiasi terbaik di wakil Kelantan untuk AR yang berlainan. Sementara itu untuk Terengganu, RCP2.6 (AR5) dan SRES B2 (AR4) telah dipilih kerana prestasi % MAE yang paling kecil. Keputusan iklim yang diunjurkan dijangka mempunyai kenaikan minimum pada maksimum (0.79%), purata (0.43%) dan suhu min (0.2%). Curah hujan setempat memperlihatkan peningkatan pola dengan (9.37%) di Stesen Gunong Barat Bachok, (5.04%) untuk Stesen Rumah Pam Salor Pengkalan Kubor, (9.11%) untuk Stesen Sg. Simpang Ampat di Kelantan. Bagi Terengganu, pola menunjukkan kenaikan (4.43%) untuk Station Sek Men. Bukit Sawa, (5.25%) untuk Stesen Rumah Pam Pulau Musang, dan (42.07%) untuk Stesen Kg Peringat.

ABSTRACT

The Intergovernmental Panel on Climate Change (IPCC) revised the impact of greenhouse gases (GHGs) into the climate system and came out with the Fifth Assessment Report (AR5) in year 2014. By AR5, the climate changes impact were classified based on the level of radiation forcing known as Representative Concentration Pathways (RCP2.6, RCP4.5, and RCP8.5). This version was slightly difference with the AR4 version which based on the GHGs groups known as A1, B1, A2, and B2. The modification in the climate changes assessment will affecting the accuracy of the climate projection in the long term. Therefore, the main aim of this study was to determine the impact in the climate change assessment between Fourth Assessment Report (AR4) with AR5. The study was focused on Eastern Malaysia including Kelantan and Terengganu. In this study, the Statistical Downscaling Model (SDSM) was used as a statistical climate model to assess the differences of the climate performances. Meanwhile, the general circulation model (GCM) provided by Climate Modelling and Analysis (CanESM2) was used for the long-term climate generation. Referring to the results, the predictor of p-u, r500, and r850 are the most influence variables in forming the local temperature and rainfall at the regions. The accuracy of the climate generation was controlled by the lower %MAE with high Correlation (R) in the calibrated and validated results. The temperature simulation was successfully to produce 0.6% of %MAE with R close to 1.0. Meanwhile the rainfall at Terengganu and Kelantan were produced less than 14% of %MAE with 0.99 of R. Based on the comparison performances between GCMs and historical data, the RCP4.5 (AR5) and SRES A2 (AR4) have been selected as the best radiation forcing level at Kelantan representative for different ARs. Meanwhile for Terengganu, the RCP2.6 (AR5) and SRES B2 (AR4) have been selected due to least %MAE performances. The projected climate results were expected to have minimum increment in the max (0.79%), mean (0.43%) and min (0.2%) temperature. The local rainfall shows increasing pattern with (9.37%) in Station Gunong Barat Bachok, (5.04%) for Station Rumah Pam Salor Pengkalan Kubor, (9.11%) for Station Sg. Simpang Ampat in Kelantan. For Terengganu, the pattern shows an increment of (4.43%) for Station Sek Men. Bukit Sawa, (5.25%) for Station Rumah Pam Pulau Musang, and (42.07%) for Station Kg Peringat.

TABLE OF CONTENT

DECLARATION	
TITLE PAGE	
ACKNOWLEDGEMENTS	ii
ABSTRAK	iii
ABSTRACT	iv
TABLE OF CONTENT	v
3LIST OF FIGURES	ix
LIST OF SYMBOLS	xi
LIST OF ABBREVIATIONS	xii
CHAPTER 1 INTRODUCTION	1
1.1 Introduction	1
1.2 Problem Statement	3
1.3 Objectives of The Study	5
1.4 Scope of Study	5
1.5 Significance of The Study	6
CHAPTER 2 LITERATURE REVIEW	7
2.1 Introduction	7
2.2 Climate Projection	9
2.2.1 Increase in Temperature	12
2.2.2 Global Warming	16
2.3 Greenhouse Gases	19

2.3.1	Components of These Gases	21
2.3.2	Climate Affected by These Models	21
2.4	General Circulation Model	23
2.4.1	Fourth Assessment Report	25
2.4.2	Fifth Assessment Report	26
2.4.3	Differences Between AR4 & AR5	27
2.4.4	General Circulation Model	29
2.5	Types of Climate Model	31
2.5.1	Regional Climate Model	31
2.5.2	Dynamic Downscaling	34
2.5.3	Statistical Downscaling	35
CHAPTER 3 METHODOLOGY		36
3.1	Introduction	36
3.2	Statistical Downscaling Model (SDSM)	37
3.3	Atmospheric Characteristics of the GCMs	40
3.4	Construction of The Climate Change Scenario	41
3.5	Calibration & Validation Process in SDSM	42
3.6	Location of Study	43
CHAPTER 4 RESULTS AND DISCUSSION		46
4.1	Introduction	46
4.2	Predictors Selection	48
4.3	Calibrated and Validated Performances	49
4.3.1	Temperature Result	49
4.3.2	Rainfall Result Kelantan	52

4.3.3	Rainfall Result Terengganu	55
4.4	Best RCP & SRES Analysis	58
4.4.1	Best RCP Analysis	58
4.4.1.1	Best RCP Temperature	58
4.4.1.2	Best RCP Rainfall Kelantan	62
4.4.1.3	Best RCP Rainfall Terengganu	66
4.4.2	Best SRES Analysis	71
4.4.2.1	Best SRES Temperature	72
4.4.2.2	Best SRES Rainfall Kelantan	75
4.4.2.3	Best SRES Rainfall Terengganu	79
4.5	Generation of Future Climate Trends of Kelantan & Terengganu using predictors AR4 & AR5	83
4.5.1	Temperature	84
4.5.2	Rainfall	87
4.5.2.1	Kelantan State	88
4.5.2.2	Terengganu State	91
CHAPTER 5 CONCLUSION		95
5.1	Conclusion	95
5.1.1	Best RCP & SRES for Terengganu & Kelantan	95
5.1.2	Performance difference between SRES and RCP	96
REFERENCES		98

LIST OF TABLES

Table 3.1	Predictor List	40
Table 4.1	List of Stations with Data, name and calibration & validation year	47
Table 4.2	List of selected predictors	48
Table 4.3	Temperature Calibration & Validation	49
Table 4.4	Rainfall Kelantan Calibration & Validation	52
Table 4.4	Rainfall Terengganu Calibration & Validation	55
Table 4.6	Temperature Best RCP Analysis	59
Table 4.7	Kelantan Rainfall Best RCP Analysis	63
Table 4.8	Terengganu Rainfall Best RCP Analysis	67
Table 4.9	Temperature Best SRES Analysis	72
Table 4.10	Kelantan Rainfall Best SRES Analysis	76
Table 4.11	Terengganu Rainfall Best SRES Analysis	80
Table 4.12	Temperature Projection	86
Table 4.13	Kelantan Rainfall Projection	90
Table 4.14	Terengganu Rainfall Projection	93

LIST OF FIGURES

Figure 2.1	SRES Storylines	10
Figure 2.2	RCP Pathways in the Fifth Assessment Report (AR5)	12
Figure 2.3	Global Surface Temperature	13
Figure 2.4	Annual Global Precipitation	15
Figure 2.5	CIMP5 Models of RCP scenarios	18
Figure 2.6	Radiative forcing elements in RCP	19
Figure 2.7	GCM concept	24
Figure 2.8	SRES emissions scenarios with the amount of GHG in every scenario	28
Figure 2.9	Graph of RCP scenarios againsts carbon dioxide emissions	29
Figure 2.10	Gridlines in GCM models	31
Figure 2.11	RCM Simulation example	32
Figure 2.12	Graph of RCM Simulation	33
Figure 2.13	Projected change in precipitation	33
Figure 3.1	Methodology of the study	37
Figure 3.2	General layout of methodology coupled with SDSM usage	38
Figure 3.3	Full SDSM Methodlogy of the downscaling method	39
Figure 3.4	Map of Kelantan	44
Figure 3.5	Map of Terengganu	45
Figure 4.1	Temperature Calibration & Validation	51
Figure 4.2	Kelantan Rainfall Calibration & Validation	54
Figure 4.2	Terengganu Rainfall Calibration & Validation	57
Figure 4.4	Temperature Best RCP:- a) Temperature vs Historical. b) Yealy Average of Temperature Vs Historical	61
Figure 4.5	Kelantan Rainfall best RCP:- a) Rainfall Monthly Average vs Historical. b) Yealy Average of Rainfall Vs Historical	65
Figure 4.6	Terengganu Rainfall Best RCP: - a) Rainfall Monthly Average vs Historical. b) Yealy Average of Rainfall Vs Historical	69
Figure 4.7	Temperature Best SRES:- a) Temperature vs Historical. b) Yealy Average of Temperature Vs Historical	74
Figure 4.8	Kelantan Rainfall Best SRES: - a) Rainfall Monthly Average vs Historical. b) Yealy Average of Rainfall Vs Historical	78
Figure 4.9	Terengganu Rainfall Best SRES: - a) Rainfall Monthly Average vs Historical. b) Yealy Average of Rainfall Vs Historical	82
Figure 4.10	Temperature Projection over 30-year intervals	85

Figure 4.11	Kelantan Rainfall Projection over 30-year intervals	89
Figure 4.12	Terengganu Rainfall Projection over 30-year intervals	92

LIST OF SYMBOLS

%	Percentage
R	Correlation value

LIST OF ABBREVIATIONS

CIMP5	Coupled Intercomparison Model ver. 5
AR5	Fifth Assessment Report
AR4	Fourth Assessment Report
SRES	Special Report on Emission Scenarios
HadCM3	Hadley Center Coupled Model
RCP	Representative Concentration Pathways
CanESM2	Canadian Center for Climate Modelling Analysis

CHAPTER 1

INTRODUCTION

1.1 Introduction

Climate is the weather conditions of an area in general or for over a long period. Climate can be assessed by long term study of the weather of a certain place. The climate can be assessed by a myriad of parameters including temperature, humidity, atmospheric pressure, wind and also precipitation. These parameters can lead us in deciding what the climate of the certain area is and in turn, from climate it was possible to gain the climate trend in which can be assess the changes of the climate towards the future. Climate trend depends on these parameters that can change involuntarily or voluntarily that depends on human actions. These trends can affect a certain area in many ways, such as the increase in temperature can lead to continued warming and many more. To assess these models, it was difficult as it requires to have a proper knowledge on how the cycle works and how are mankind going to interpret the data from General Circulation Model (GCM).

Climate is commonly defined as the weather over a long period. The standard averaging period was 30 years. The period can be lengthened or shorten the period depending on the purpose of the study. It also includes statistics, other than daily averages, such as the magnitudes of day-to-day or year to year variations. In climate, there stands a term called climate normal. These terms were actually a reference point for the climatologists to identify what went astray from the normal climate. A climate that follows the pattern of recent and past climates was a normal climate. In the span of 30 years used as the period of the study, if there are any unnatural occurrences of weather extremes such as heat waves, or heavy precipitation, it can be recorded into the journal for reference about the climate trend, and where the climate trend was heading.

Climate projection is another branch in climate change. Climate projection much more focuses on projections of the climate towards the future. Climate projections are mainly based on the current climate and the climate trend that was ongoing. Climate projection can be also defined as the stimulated response of the climate system to the scenario of future emission or concentration of greenhouse gases and aerosols, that are generally derived using climate models. Climate projections are distinguished from climate predictions by their dependence on the emission, the concentration and the radiative forcing scenario used, which was based on assumptions concerning future socioeconomic and technological developments that may or may not be realized. Climate projection has been widely used since the early twentieth century where it is viable to predict the climate change by having calculated the emissions of gases in the atmosphere. Projection can also be determined by the increment of temperature of the atmosphere, whilst neglecting to project climate for the future, it can lead to several problems including decreased productivity in agricultural lands, submerging of suburban areas due to floods or heavy precipitation.

The scientific community has reached a 97% consensus that climate change is influenced by humans yet many people still doubt that this was true (Thompson, 2017). The problems that are in climate can be catastrophic if left unattended. The major climate problems are that there is no correct way of measuring projection in terms of climate. The climate itself presents a question to researchers. The climate itself was handled by the IPCC or the Intergovernmental Panel on Climate Change which was responsible for the climate projection in which the researchers, working for the IPCC assess the climate trend and calculate the climatic change to obtain the projected climate in the future.

The problem arises when there are inconsistencies in certain climatic trends in which can contribute to projection problems. Not forgetting, global climate problems also pave the way to the obstacle in generating climate projections. Such problems, including unnatural climate trends, caused by the presence of GHG in the stratosphere have curbed researchers advances in climate prediction. With the implementation of the Assessment Reports, specifically the Fourth Assessment Report, a system, namely SRES or the Special Reports on Emission Concentration was used to assess the climatic changes and

the climatic problems thus being able to project the climate of the world. The system, is a cry away from perfectness, as the system overestimates the variables that are the guidelines to climate projection. With the implementation of the Fifth Assessment Report, a new system was constructed and it supersedes the SRES. The system, namely RCP focuses on the endpoints and up until now it has been accepted as a sound system by the IPCC. The time to time revision of the climate assessment is necessary due to several issues such as the policy-relevant calculation and overestimate responses of the GHG emissions forcing, reducing regional biases in temperature simulation, the correlation between observed and modelled mean precipitation, uncertainties of cloud processes (Tukimat *et. al.*, 2014).

1.2 Problem Statement

To the International Panel on Climate Change (IPCC) has assessed the amounts of greenhouse gases GHG in various areas of the world. And from these greenhouse gases, the researchers adopted four pathways that are the most prevalent in the stratosphere in today's community. There was a clear view on the increase of the Representative Concentration Pathways (RCPs) throughout the years. The four pathways that have been selected are used for climate modelling and research, which all of them describe different climate projection, all of these pathways consider the amount of GHGs that are emitted in the years to come. The four RCPs, namely RCP2.6, RCP4.5, RCP6 and RCP8.5 are labelled according to possible range of radiative forcing values in the year 2100 in relative to pre-industrial values.

Also assessed by the council of IPCC, the Fourth Assessment Report uses a system that is different from the system that was used in AR5. with the usage of SRES in AR4, there are quite many scenarios that can be obtained through these scenarios. SRES scenarios are emission scenarios developed by (Nakicenovic and Swart, 2000) and used, among others, as a basis for some of the climate projections used in the Fourth Assessment Report. This assessment system was obsolete and was superseded by the Fifth Assessment Report's system which uses the RCP or the Representative Concentration Pathways system. Still, both systems measure and record the increase of the values of GHG and the increase of other features and indicators that are used in measuring the effect of the GHG .

REFERENCES

- Alves, J. M. B., Junior, F. C. V., Chaves, R. R., Silva, E. M., Servain, J., Costa, A. A., ... Acb, B. (2016). Evaluation of the AR4 CMIP3 and the AR5 CMIP5 Model and Projections for Precipitation in Northeast Brazil, 4(April). <https://doi.org/10.3389/feart.2016.00044>
- Ayugi, B. O., Wang, W., & Chepkemai, D. (2016). Analysis of Spatial and Temporal Patterns of Rainfall Variations over Kenya. *Environment and Earth Science*, 6(11), 69–83.
- Chen, S., Yu, P., & Tang, Y. (2010). Statistical downscaling of daily precipitation using support vector machines and multivariate analysis. *Journal of Hydrology*, 385(1–4), 13–22. <https://doi.org/10.1016/j.jhydrol.2010.01.021>
- Faiz, M. A., Liu, D., Fu, Q., Li, M., Baig, F., Tahir, A. A., ... Cui, S. (2018). Performance evaluation of hydrological models using ensemble of General Circulation Models in the northeastern China. *Journal of Hydrology*, 565(August), 599–613. <https://doi.org/10.1016/j.jhydrol.2018.08.057>
- Giorgi, F. (2010). Uncertainties in climate change projections, from the global to the regional scale. *EPJ Web of Conferences*, 9, 115–129. <https://doi.org/10.1051/epjconf/201009009>
- Hannah, L. (2015). *The Climate System and Climate Change. Climate Change Biology*. <https://doi.org/10.1016/B978-0-12-420218-4.00002-0>
- IPCC. (2014). *Climate Change 2014: Synthesis Report. Climate Change 2014: Synthesis*. <https://doi.org/10.1256/004316502320517344>
- Jalota, S. K., Vashisht, B. B., Sharma, S., & Kaur, S. (2018). *Climate Change Projections. Understanding Climate Change Impacts on Crop Productivity and Water Balance*. <https://doi.org/10.1016/B978-0-12-809520-1.00002-1>
- Jian-Bin, H., Shao-Wu, W., Yong, L., Zong-Ci, Z., & Xin-Yu, W. (2012a). Debates on the Causes of Global Warming. *Advances in Climate Change Research*, 3(1), 38–44. <https://doi.org/10.3724/SP.J.1248.2012.00038>
- Jian-Bin, H., Shao-Wu, W., Yong, L., Zong-Ci, Z., & Xin-Yu, W. (2012b). The Science of Global Warming. *Advances in Climate Change Research*, 3(3), 174–178. <https://doi.org/10.3724/SP.J.1248.2012.00174>
- Le Roux, R., Katurji, M., Zawar-Reza, P., Quénol, H., & Sturman, A. (2018). Comparison of statistical and dynamical downscaling results from the WRF model. *Environmental Modelling and Software*, 100, 67–73. <https://doi.org/10.1016/j.envsoft.2017.11.002>
- Nadrah, N., Tukimat, A., Shahid, S., Isabella, J., & Gisen, A. (2014). ASSESSING THE PERFORMANCES OF THE GLOBAL CLIMATE MODELS (AR4 & AR5

) IN LONG-TERM PROJECTION OF CLIMATE IN MALAYSIA, 5.

- Najafi, M. R., Asce, S. M., Moradkhani, H., Asce, M., Wherry, S. A., & Asce, S. M. (2011). Statistical Downscaling of Precipitation Using Machine Learning with Optimal Predictor Selection, (August), 650–664.
[https://doi.org/10.1061/\(ASCE\)HE.1943-5584.0000355](https://doi.org/10.1061/(ASCE)HE.1943-5584.0000355).
- Sachindra, D. A., Huang, F., Barton, A., & Perera, B. J. C. (2013). Least square support vector and multi-linear regression for statistically downscaling general circulation model outputs to catchment streamflows, *1106*(April 2012), 1087–1106.
<https://doi.org/10.1002/joc.3493>
- Sachindra, D. A., Huang, F., Barton, A., & Perera, B. J. C. (2014). Statistical downscaling of general circulation model outputs to precipitation – part 2 : bias-correction and future, *3303*(January), 3282–3303. <https://doi.org/10.1002/joc.3915>
- Schwartz, S. E. (2018). *Unrealized Global Temperature Increase: Implications of Current Uncertainties*. *Journal of Geophysical Research: Atmospheres* (Vol. 123).
<https://doi.org/10.1002/2017JD028121>
- Sidharno, W. (2016). STUDY of CLIMATE CHANGE IMPACT on WATER AVAILABILITY in OEBOBO LILIBA WATERSHED, (July), 29–31.
- Thompson, J. E. (2017). Survey data reflecting popular opinions of the causes and mitigation of climate change. *Data in Brief*, *14*, 412–439.
<https://doi.org/10.1016/j.dib.2017.07.060>
- Tukimat, N. N. A., & Alias, N. A. (2016). Assessment the potential of SRES scenario for Kuala Sala, Malaysia. *J. Mech. Civ. Eng*, *13*(3), 6–12.
<https://doi.org/10.9790/1684-1303010612>
- Vargas, X., Gómez, T., Ahumada, F., Rubio, E., Cartes, M., & Gibbs, M. (2013). Water availability in a mountainous Andean watershed under CMIP5 climate change scenarios, *2013*(July), 33–38.
- Walz, A., Braendle, J. M., Lang, D. J., Brand, F., Briner, S., Elkin, C., ... Schmatz, D. R. (2014). Experience from downscaling IPCC-SRES scenarios to specific national-level focus scenarios for ecosystem service management. *Technological Forecasting and Social Change*, *86*, 21–32.
<https://doi.org/10.1016/j.techfore.2013.08.014>